

Swine Flu & YOU



What is swine flu?

Swine Influenza (swine flu) is a respiratory disease of pigs caused by type A influenza viruses that causes regular outbreaks in pigs. People do not normally get swine flu, but human infections can and do happen.

Is this swine flu virus contagious?

CDC has determined that this swine influenza A (H1N1) virus is contagious and is spreading from human to human. However, at this time, it is not known how easily the virus spreads between people.

What are the signs and symptoms of swine flu in people?

The symptoms of swine flu in people are similar to the symptoms of regular human flu and include fever, cough, sore throat, body aches, headache, chills and fatigue. Some people have reported diarrhea and vomiting associated with swine flu. In the past, severe illness (pneumonia and respiratory failure) and deaths have been reported with swine flu infection in people. Like seasonal flu, swine flu may cause a worsening of underlying chronic medical conditions.

How does swine flu spread?

Spread of this swine influenza A (H1N1) virus is thought to be happening in the same way that seasonal flu spreads. Flu viruses are spread mainly from person to person through coughing or sneezing of people with influenza. Sometimes people may become infected by touching something with flu viruses on it and then touching their mouth or nose.

How can someone with the flu infect someone else?


Infected people may be able to infect others beginning 1 day before symptoms develop and up to 7 or more days after becoming sick. That means that you may be able to pass on the flu to someone else before you know you are sick, as well as while you are sick.

Are there medicines to treat swine flu?

Yes. CDC recommends the use of oseltamivir or zanamivir for the treatment and/or prevention of infection with these swine influenza viruses. Antiviral drugs are prescription medicines (pills, liquid or an inhaler) that fight against the flu by keeping flu viruses from reproducing in your body. If you get sick, antiviral drugs can make your illness milder and make you feel better faster. They may also prevent serious flu complications. For treatment, antiviral drugs work best if started soon after getting sick (within 2 days of symptoms).

What surfaces are most likely to be sources of contamination?

Germs can be spread when a person touches something that is contaminated with germs and then touches his or her eyes, nose, or mouth. Droplets from a cough or sneeze of an infected person move through the air. Germs can be spread when a person touches respiratory droplets from another person on a surface like a desk and then touches their own eyes, mouth or nose before washing their hands.



Can I get swine flu from eating or preparing pork?

No. Swine influenza viruses are not spread by food. You cannot get swine influenza from eating pork or pork products. Eating properly handled and cooked pork products is safe.

How long can viruses live outside the body?

We know that some viruses and bacteria can live 2 hours or longer on surfaces like cafeteria tables, doorknobs, and desks. Frequent handwashing will help you reduce the chance of getting contamination from these common surfaces.

What should I do if I get sick?

If you live in areas where swine influenza cases have been identified and become ill with influenza-like symptoms, including fever, body aches, runny nose, sore throat, nausea, or vomiting or diarrhea, you may want to contact their health care provider, particularly if you are worried about your symptoms. Your health care provider will determine whether influenza testing or treatment is needed.

If you are sick, you should stay home and avoid contact with other people as much as possible to keep from spreading your illness to others.

If you become ill and experience any of the following warning signs, seek emergency medical care.

In children emergency warning signs that need urgent medical attention include:

- » **Fast breathing or trouble breathing**
- » **Bluish skin color**
- » **Not drinking enough fluids**
- » **Not waking up or not interacting**
- » **Being so irritable that the child does not want to be held**
- » **Flu-like symptoms improve but then return with fever and worse cough**
- » **Fever with a rash**
- » **In adults, emergency warning signs that need urgent medical attention include:**
- » **Difficulty breathing or shortness of breath**
- » **Pain or pressure in the chest or abdomen**
- » **Sudden dizziness**
- » **Confusion**
- » **Severe or persistent vomiting**

Content source: Swine Influenza and You, Centers for Disease Control and Prevention

http://www.cdc.gov/swineflu/swineflu_you.htm

Please contact Symmetry Wellness for support or training on this topic.